

TOPIC 20 EXERCISE 3 – PROTON NMR SPECTRA

1. A compound with molecular formula $C_3H_6O_2$ gives the following peaks in its proton nmr spectrum:

| Chemical shift | Splitting | Integration factor |
|----------------|-----------|--------------------|
| 1.1 | Triplet | 3 |
| 2.2 | Quartet | 2 |
| 11.8 | Singlet | 1 |

Identify the molecule and account for the chemical shifts, splitting and integration factors of all three peaks.


2. A compound with molecular formula $C_5H_{10}O_2$ gives the following peaks in its proton nmr spectrum:

| Chemical shift | Splitting | Integration factor |
|----------------|-----------|--------------------|
| 1.2 | Triplet | 3 |
| 1.3 | Triplet | 3 |
| 2.3 | Quartet | 2 |
| 4.1 | Quartet | 2 |

- a) Identify the molecule and account for the chemical shifts, splitting and integration factors of all four peaks.
- b) Explain why CH_3Cl is not used as a solvent in proton nmr spectroscopy.
- c) Give three reasons why T.M.S. is a good standard in proton nmr spectroscopy.
3. Use the information in the table below to identify molecule A from its proton nmr spectrum:


| peak | chemical shift/ppm | relative intensity | peak type |
|------|--------------------|--------------------|-----------|
| a | 2.4 | 1 | multiplet |
| b | 2.1 | 3 | Singlet |
| c | 1.5 | 2 | multiplet |
| d | 1.1 | 3 | doublet |
| e | 0.9 | 3 | triplet |

4. Identify the molecule responsible for the proton nmr spectrum below:




The peak at 9.6 is a doublet; the peak at 2.1 is a multiplet; the peak at 1.7 is a multiplet and the peak at 0.9 is a triplet

5. Identify this molecule:




| Shift | Relative peak intensity | Splitting |
|-------|-------------------------|-----------|
| 4.1 | 2 | Quartet |
| 2.1 | 3 | Singlet |
| 1.3 | 3 | Triplet |

6. Identify this molecule:




| Shift | Relative peak intensity | Splitting |
|-------|-------------------------|-----------|
| 2.4 | 2 | Quartet |
| 2.2 | 3 | Singlet |
| 1.1 | 3 | Triplet |

7. Identify this molecule:



| Shift | Relative peak intensity | Splitting |
|-------|-------------------------|-----------|
| 3.6 | 1 | Multiplet |
| 2.3 | 1 | Singlet |
| 1.4 | 2 | Multiplet |
| 1.2 | 3 | Doublet |
| 0.9 | 3 | Triplet |


8. Identify this molecule:



HSP-00-115

| Shift | Relative peak intensity | Splitting |
|-------|-------------------------|-----------|
| 11.6 | 1 | Singlet |
| 2.2 | 2 | Triplet |
| 1.9 | 2 | Multiplet |
| 1.0 | 3 | Triplet |


9. Identify this molecule:



| Shift | Relative peak intensity | Splitting |
|-------|-------------------------|-----------|
| 9.6 | 1 | Doublet |
| 2.3 | 1 | Multiplet |
| 1.1 | 6 | Doublet |


SOLUTIONS

- peak at 1.1 is CH_3 - adjacent to $-\text{CH}_2$ -
peak at 2.2 is $-\text{CH}_2\text{CO}$ -, adjacent to CH_3 -
peak at 11.8 is $-\text{COOH}$
so molecule is propanoic acid, $\text{CH}_3\text{CH}_2\text{COOH}$
- peak at 1.2 is CH_3 -, adjacent to $-\text{CH}_2$ -
peak at 1.3 is also CH_3 -, adjacent to $-\text{CH}_2$ -
peak at 2.3 is $-\text{CH}_2\text{CO}$ -, adjacent to CH_3 -
peak at 4.1 is $-\text{CH}_2\text{O}$ -, adjacent to CH_3 -
so molecule is ethyl propanoate, $\text{CH}_3\text{CH}_2\text{COOCH}_2\text{CH}_3$
 - CHCl_3 is not used as a solvent because it contains a proton which will interfere with the spectrum of the substance being analysed.
 - TMS is a good standard because
 - it contains 12 identical protons, giving a single intense peak
 - it contains highly shielded protons, which do not interfere with the spectrum
 - it is cheap and non-toxic
- e is $-\underline{\text{CH}}_3$ adjacent to $-\underline{\text{CH}}_2$ - (c)
b is $-\underline{\text{CH}}_3$ adjacent to $-\text{C}=\text{O}$
d is $-\underline{\text{CH}}_3$ adjacent to $-\underline{\text{CH}}-$ (a)
a is $-\underline{\text{CH}}-$ adjacent to $-\text{C}=\text{O}$



3-methylpentan-2-one

- 9.6 is $-\underline{\text{C}}\text{HO}$ adjacent to $-\underline{\text{C}}\text{H}$ - (2.1)
2.1 is $-\underline{\text{C}}\text{H}$ - adjacent to $-\underline{\text{C}}\text{HO}$ (9.6)
0.9 is 2 x $-\underline{\text{C}}\text{H}_3$ adjacent to $-\underline{\text{C}}\text{H}_2$ - (1.7)



5. 4.1 is O-CH₂- adjacent to -CH₃ (1.3)
 2.1 is CH₃C=O
CH3COOCH2CH3 (ethyl ethanoate)
 (2.1) (4.1)(1.3)
6. 2.4 is O=CCH₂ adjacent to CH₃ (1.1)
 2.2 is CH₃C=O
CH3COCH2CH3 (butanone)
 (2.2) (2.4)(1.1)
7. 0.9 is CH₃ adjacent to CH₂ (1.4)
 1.2 is CH₃ adjacent to CH (3.6)
 3.6 is O-CH-
 2.3 is -OH
CH3CH2CH(OH)CH3 (butan-2-ol)
 (0.9)(1.4)(3.6)(2.3)(1.2)
8. 11.6 is -COOH
 2.2 is O=CCH₂- adjacent to -CH₂- (1.9)
 1.0 is -CH₃ adjacent to -CH₂- (1.9)
CH3CH2CH2COOH (butanoic acid)
 (1.0)(1.9)(2.2) (11.6)
9. 9.6 is -CHO adjacent to -CH- (2.3)
 2.3 is -CHC=O
 1.1 is 2 x CH₃ adjacent to -CH- (2.3)
(CH3)2CHCHO (methylpropanal)
 (1.1) (2.3)(9.6)